

CORONAVIRUS

SARS-CoV-2 seroprevalence among parturient women in Philadelphia

Dustin D. Flannery^{1,2,3*}, Sigrid Gouma^{4*}, Miren B. Dhudasia^{1,3}, Sagori Mukhopadhyay^{1,2,3}, Madeline R. Pfeifer¹, Emily C. Woodford¹, Jeffrey S. Gerber^{2,3,5}, Claudia P. Arevalo⁴, Marcus J. Bolton⁴, Madison E. Weirick⁴, Eileen C. Goodwin⁴, Elizabeth M. Anderson⁴, Allison R. Greenplate^{6,7}, Justin Kim^{6,7}, Nicholas Han^{6,7}, Ajinkya Pattekar^{6,8}, Jeanette Dougherty^{6,7}, Oliva Kuthuru^{6,7}, Divij Mathew^{6,7}, Amy E. Baxter^{6,7}, Laura A. Vella^{5,6}, JoEllen Weaver⁹, Anurag Verma¹⁰, Rita Leite¹¹, Jeffrey S. Morris¹², Daniel J. Rader^{9,10}, Michal A. Elovitz^{6,11}, E. John Wherry^{6,7}, Karen M. Puopolo^{1,2,3†}, Scott E. Hensley^{4,6†}

Limited data are available for pregnant women affected by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). Serological tests are critically important for determining SARS-CoV-2 exposures within both individuals and populations. We validated a SARS-CoV-2 spike receptor binding domain serological test using 834 pre-pandemic samples and 31 samples from COVID-19–recovered donors. We then completed SARS-CoV-2 serological testing of 1293 parturient women at two centers in Philadelphia from 4 April to 3 June 2020. We found 80 of 1293 (6.2%) of parturient women had immunoglobulin G (IgG) and/or IgM SARS-CoV-2–specific antibodies. We found race/ethnicity differences in seroprevalence rates, with higher rates in Black/non-Hispanic and Hispanic/Latino women. Of the 72 seropositive women who also received nasopharyngeal polymerase chain reaction testing during pregnancy, 46 (64%) were positive. Continued serologic surveillance among pregnant women may inform perinatal clinical practices and can potentially be used to estimate exposure to SARS-CoV-2 within the community.

INTRODUCTION

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) can cause serious disease in adult populations, particularly in those with underlying health conditions (1). Serological tests are important for determining SARS-CoV-2 viral exposures within individuals and populations (2). However, many commercial tests have high false-positive rates and therefore cannot be used to accurately estimate seroprevalence in populations with relatively low levels of exposures (3, 4). Serological tests are especially important for vulnerable populations such as pregnant women, because immune status has implications for management of both the pregnant woman and the newborn. Admission to the hospital for delivery is one of the few instances in which otherwise healthy individuals are consistently interacting with the medical system and therefore provides an opportunity for surveillance of SARS-CoV-2 serology in the community.

¹Division of Neonatology, Children's Hospital of Philadelphia, Philadelphia, PA, USA. ²Department of Pediatrics, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA, USA. ³Center for Pediatric Clinical Effectiveness, Children's Hospital of Philadelphia, Philadelphia, PA, USA. ⁴Department of Microbiology, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA, USA. ⁵Division of Infectious Diseases, Children's Hospital of Philadelphia, Philadelphia, PA, USA. ⁶Institute for Immunology, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA, USA. ⁷Department of Systems Pharmacology and Translational Therapeutics, University of Pennsylvania, Philadelphia, PA. ⁸Division of Gastroenterology, Department of Medicine, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA, USA. ⁹Institute for Translational Medicine and Therapeutics, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA, USA. ¹⁰Departments of Genetics and Medicine, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA. ¹¹Maternal and Child Health Research Center, Department of Obstetrics and Gynecology, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA, USA. ¹²Department of Biostatistics Epidemiology and Informatics, University of Pennsylvania, Philadelphia, PA, USA. *These authors contributed equally to this work.

†Corresponding author. Email: hensley@penmedicine.upenn.edu (S.E.H.); karen.puopolo@penmedicine.upenn.edu (K.M.P.)

We performed a study of pregnant women presenting for delivery from 4 April to 3 June 2020 at two academic birth hospitals in Philadelphia, Pennsylvania. Both hospitals are active clinical and research centers affiliated with the University of Pennsylvania and, when combined, represent 50% of live births in Philadelphia (5). Discarded maternal sera from delivery admission were collected, de-identified, and tested by enzyme-linked immunosorbent assay (ELISA) for SARS-CoV-2 immunoglobulin G (IgG) and IgM antibodies to the spike receptor binding domain (RBD) antigen.

RESULTS

Demographics

Demographics and clinical characteristics of the women are shown in Table 1. Most serum specimens were derived from women living in areas within or immediately bordering the city of Philadelphia (Fig. 1). Pregnant women who were symptomatic or exposed to SARS-CoV-2 underwent SARS-CoV-2 nasopharyngeal nucleic acid polymerase chain reaction (PCR) testing from 4 to 12 April 2020; universal PCR testing was recommended for all pregnant women presenting for delivery starting 13 April 2020. Of 1514 women who delivered during the study period, 1293 (85%) had available discarded serum specimens and were included in the analysis.

Assay validation

Our serological assay used a SARS-CoV-2 spike RBD antigen and modified ELISA protocol first described by Amanat *et al.* (6). We validated this serological assay by testing serum samples collected before the pandemic in 2019 from 834 individuals in the Penn Medicine BioBank (PMBB) and 31 individuals who recovered from confirmed coronavirus disease 19 (COVID-19) infections in 2020 (Fig. 2, A and B). All 31 serum samples from COVID-19–recovered donors contained high, but variable, levels of SARS-CoV-2 IgG (Fig. 2A), and

Table 1. Demographics and clinical characteristics of the study cohort. IQR, interquartile range; BMI, body mass index.

Characteristics	Total (n = 1293)	Seropositive* (n = 80)	Seronegative (n = 1213)	P value [†]
Age (in years), median (IQR)	31 (27, 35)	28 (24, 32)	31 (27, 35)	<0.001
Race/ethnicity, n (%)[‡]				
Black/non-Hispanic	537	52 (9.7)	485 (90.3)	<0.001
White/non-Hispanic	447	9 (2.0)	438 (98.0)	<0.001
Hispanic/Latino	125	13 (10.4)	112 (89.6)	0.04
Asian	106	1 (0.9)	105 (99.1)	0.01
Other/unknown [§]	78	5 (6.4)	73 (93.6)	0.93
Pre-pregnancy BMI, n (%)[‡]				
Overweight (25.0 to <30.0)	345	28 (8.1)	317 (91.9)	0.07
Obese (≥30.0)	337	27 (8.0)	310 (92.0)	0.09
Diabetes[¶], n (%)[‡]	113	10 (8.9)	103 (91.1)	0.22
Hypertension[¶], n (%)[‡]	404	33 (8.2)	371 (91.8)	0.05
Asthma[¶], n (%)[‡]	194	13 (6.7)	181 (93.3)	0.75
Cesarean delivery, n (%)[‡]	400	30 (7.5)	370 (92.5)	0.19
Preterm delivery at gestational age <37 weeks, n (%)[‡]	128	11 (8.6)	117 (91.4)	0.23
Live-born infant, n (%)[‡]	1282	79 (6.2)	1203 (93.8)	0.51

*Seropositivity was based on either IgG or IgM level of >0.48 arbitrary units. †Difference in maternal age was tested using Mann-Whitney *U* test, differences in proportion of all other characteristics were tested using χ^2 tests or Fisher's exact test as appropriate. For race/ethnicity and pre-pregnancy BMI, difference was tested at each level of the characteristic (e.g., proportion of Black women who were seropositive compared with proportion of non-Black women who were seropositive). ‡Row percentages are shown, which represent the percent of total in each characteristic (e.g., 9.7% of Black/Non-Hispanic women were seropositive). §Race/ethnicity was unknown for two seropositive and 26 seronegative women; race was abstracted from documentation at the time of admission and in clinical practice and is usually self-reported. ||Pre-pregnancy BMI was missing for 2 seropositive and 12 seronegative women; pre-pregnancy BMI was abstracted from documentation in the medical record or from patient's self-reported entry in birth registration. ¶Diagnoses were based on delivery admission ICD-10 diagnosis codes for diabetes (O24, E08-E13, and Z79.4), hypertension (O10, O11, O13-O16, I10-I13, and I15), and asthma (J45).




Fig. 1. Geographical distribution of women tested for SARS-CoV-2 antibodies. Most serum specimens analyzed were from women living in areas within or immediately bordering the city of Philadelphia. Location of birth hospitals where serum samples were collected are shown as red crosses.

22 of 31 samples contained detectable levels of SARS-CoV-2 IgM (Fig. 2B). Conversely, only 5 of 834 samples collected before the pandemic contained SARS-CoV-2 IgG, and only 4 of 834 samples contained SARS-CoV-2 IgM; none contained both IgG and IgM. On the basis of these data, the estimated sensitivity of the test is 100%

[95% confidence interval (CI), 89.1 to 100.0%], and the specificity is 98.9% (95% CI, 98.0 to 99.5%). Using this test, we have found that there is heterogeneity in antibody responses among hospitalized COVID-19 patients and that some actively infected patients are seronegative (7, 8). Consistent with our initial validation experiments, only 1 of 140 samples collected from pregnant women before the pandemic (from 2009 to 2012) had IgG or IgM SARS-CoV-2 antibodies (Fig. 2, C and D).

Serological findings

We found that 80 of 1293 (seropositivity rate of 6.2%; 95% CI, 4.9 to 8.0%) pregnant women presenting for delivery from 4 April to 3 June 2020 had IgG and/or IgM SARS-CoV-2 antibodies (*P* = 0.003 comparing samples from pre-pandemic and pandemic pregnant women; Fig. 2, C and D). We identified 55 women with both SARS-CoV-2 IgG and IgM, 21 women with only SARS-CoV-2 IgG, and 4 women with only SARS-CoV-2 IgM (table S1). SARS-CoV-2 antibody levels in samples from these women were variable (Fig. 2, C and D), similar to what we found in samples from individuals recovering from confirmed SARS-CoV-2 infections (Fig. 2, A and B). The seroprevalence rate was not statistically different comparing women living within the city limits of Philadelphia [62 of 986 (6.3%); 95% CI, 4.9 to 8.0%] with those living in surrounding areas in Pennsylvania [12 of 191 (6.3%); 95% CI, 3.3 to 10.7%], or surrounding areas in New Jersey [5 of 107 (4.7%); 95% CI, 1.5 to 10.6%].




Fig. 2. Serum SARS-CoV-2 antibody levels in COVID-19 pandemic and pre-pandemic individuals. (A and B) Relative levels of SARS-CoV-2 IgG (A) and IgM (B) in serum collected before the COVID-19 pandemic ($n = 834$) and serum collected from COVID-19 recovered donors ($n = 31$). (C and D) Relative levels of SARS-CoV-2 IgG (C) and IgM (D) in serum collected from pregnant women from 2009 to 2012 ($n = 140$) and from 4 April to 3 June 2020 ($n = 1293$). Dashed lines indicate 0.48 arbitrary units, which was used to distinguish positive versus negative samples (see Materials and Methods). Serum samples that were below the cutoff for seropositivity were assigned an antibody level of 0.40 arbitrary units.

There were no significant differences in seroprevalence rates in women with or without comorbidities, preterm delivery, or cesarean mode of delivery (Table 1). In contrast, we observed significant race/ethnicity differences in seroprevalence rates, with higher rates in Black/non-Hispanic (9.7%; 95% CI, 7.3 to 12.5%) and Hispanic/Latino (10.4%; 95% CI, 5.7 to 17.1%) women and lower rates in White/non-Hispanic (2.0%; 95% CI, 0.9 to 3.8%) and Asian (0.9%; 95% CI, 0.0 to 5.1%) women (Table 1).

Nasopharyngeal swabs from 1109 (85.8%) women were tested by SARS-CoV-2 PCR during the pregnancy or at the time of delivery. Most of the sera tested for antibody were obtained before or within 6 days of PCR testing (Table 2). SARS-CoV-2 antibodies were detected in all sera obtained from PCR-positive women when serum samples were obtained more than 7 days after PCR testing (Table 2). Overall, we found that 46 of 72 seropositive women who were tested by PCR had a SARS-CoV-2–positive PCR result, whereas only 18 of 1037 seronegative women who were tested by PCR had a SARS-CoV-2–positive PCR result. While all serum samples were collected during the delivery admission, nasopharyngeal samples were collected at variable times either during the delivery admission or earlier in the pregnancy, and therefore, further study will be required to evaluate the temporal relationship between SARS-CoV-2 seropositivity and PCR positivity in pregnant women.

DISCUSSION

Large-scale serology testing is critical for estimating how many individuals have been infected during the COVID-19 pandemic. Because of widely imposed social distancing requirements, and to decreases in on-site, discretionary medical care, it is currently diffi-

cult to collect serum for population-wide serological testing. The vast majority of pregnant women, however, continue to have multiple interactions with the medical system for prenatal care and for delivery during this pandemic and therefore present an opportunity to consistently assess SARS-CoV-2 exposures within a community. Our data suggest that 6.2% of parturient women in Philadelphia from 4 April to 3 June 2020 were previously exposed to SARS-CoV-2.

As of 3 June 2020, there were 23,160 confirmed cases of COVID-19 in the city of Philadelphia (9), which has a population size of nearly 1.6 million people. This suggests an infection rate of about 1.4%, which is more than four times lower than the estimates based on our serological data. Serologic studies may provide a more accurate means of assessing population exposure to SARS-CoV-2 by identifying asymptomatic or minimally symptomatic and symptomatic infections. Further studies are needed to determine how the immune status of pregnant women compares with that of the general population. For example, parturient women may not represent individuals of different ages within the general population, and women and men might mount different antibody responses upon infection with SARS-CoV-2 (10). Furthermore, most pregnant women cannot fully shelter-in-place during a pandemic, as they continue to have interactions with the medical system. Our finding that Black/non-Hispanic and Hispanic/Latino women have higher SARS-CoV-2 seroprevalence rates relative to women of other races suggests that there are race/ethnicity differences in SARS-CoV-2 exposures in Philadelphia and surrounding areas. Identification of factors that contribute to such differences in exposure to SARS-CoV-2, including factors rooted in systemic racism, may inform public health measures aimed at preventing further infections (11–13).

Prior perinatal COVID-19 studies have primarily focused on virus detection (i.e., nucleic acid testing) in pregnant women, and most of these studies have not evaluated antibody responses (14–22). Two published studies to date have assessed SARS-CoV-2 serology in pregnant women with active disease. A study of six parturient women in Wuhan, China with confirmed COVID-19 found that all six women had elevated levels of SARS-CoV-2 IgG and IgM (23). A case report from Peru detailed a symptomatic pregnant woman with positive PCR testing and negative serology at presentation, who developed severe respiratory failure necessitating delivery; her IgM and IgG turned positive 4 days after delivery (9 days after symptom onset) (24). Beyond describing individual response to infection, SARS-CoV-2 serological testing among pregnant women will be increasingly important for perinatal disease risk management and for optimizing vaccine strategies when vaccines become available. Additional studies will be needed to address the impact of maternal infection on neonatal immune responses and to determine those factors that may contribute to observed disparities in exposure to SARS-CoV-2.

MATERIALS AND METHODS

Study design

The goal of this study was to estimate SARS-CoV-2 seroprevalence rates in the community using discarded serum samples from parturient women. The Institutional Review Board at the University of Pennsylvania approved this study. There was a waiver of consent for testing of residual serum samples from parturient women, as indicated below. For other cohorts used to validate our assay, subjects were consented before samples were obtained. De-identified data were used for analysis.

Table 2. Timing of serology testing and seropositivity with respect to nasopharyngeal PCR testing. The table includes 1109 women tested for serology who were also tested by nasopharyngeal PCR anytime during pregnancy up to discharge from delivery admission. NP, nasopharyngeal.

Serology timing	NP-PCR positive		NP-PCR negative	
	Tested	Seropositive (%)	Tested	Seropositive (%)
Before NP-PCR test	17	10 (58.8)	364	9 (2.5)
0–6 days after NP-PCR test	26	15 (57.7)	647	16 (2.5)
7–13 days after NP-PCR test	5	5 (100.0)	8	0
14–20 days after NP-PCR test	2	2 (100.0)	7	0
≥21 days after NP-PCR test	14	14 (100.0)	19	1 (5.3)
Total	64	46 (71.9)	1045	26 (2.5)

Serum samples from parturient women

Pregnant women at the two hospitals (Pennsylvania Hospital and Hospital of the University of Pennsylvania) have blood drawn for rapid plasma reagin (screening for syphilis per public health guidelines) testing as part of routine clinical care on admission to the hospital for delivery. Residual serum from this testing was obtained from the clinical laboratory at the time it was otherwise to be discarded. Demographic and clinical data were collected from review of electronic medical records to assess for differences in seroprevalence based on these factors. Race and ethnicity were self-reported. International Classification of Diseases, 10th revision (ICD-10) diagnosis codes O24, E08-E13, and Z79.4 were used to capture type 1 diabetes, type 2 diabetes, and gestational diabetes; codes O10, O11, O13-O16, I10-I13, and I15 were used to capture hypertensive disorders, gestational hypertension, and pre-eclampsia; and code J45 was used to capture any history of asthma before or during pregnancy. To ensure that these codes correctly captured patient condition, we manually reviewed records for the first 130 (10%) women with ICD-10 diagnosis of diabetes, hypertension, or asthma and reviewed a random sampling of 65 records from the first 130 (5% of total) women without any identified ICD-10 codes for these conditions. Patient numbers in table S1 were assigned at random. The Institutional Review Board at the University of Pennsylvania approved this study with waiver of consent.

Serum samples from individuals recovered from COVID-19

Samples from subjects who had recovered from laboratory-confirmed SARS-CoV-2 infections were obtained at the University of Pennsylvania. Subjects were consented, and samples were obtained after laboratory-confirmed COVID-19 diagnosis and >14 days since resolution of symptoms. The Institutional Review Board at the University of Pennsylvania approved this study.

Pre-pandemic human serum samples

To validate our serological assay, serum samples from 834 adults (19 to 89 years old; 52% females) were collected via the PMBB between October and December 2019, before the COVID-19 pandemic. PMBB routinely consents individuals visiting the University of Pennsylvania health care system and obtains and stores biospecimens. Banked serum samples obtained from pregnant women from 2009 to 2012 were also used as pre-pandemic controls. For these banked samples, maternal serum was collected during the third trimester of pregnancy as part of an Institutional Review Board–approved study.

From this study, 140 samples were randomly selected from women who delivered at term (average gestational age at time of sample collection was 33.8 weeks, 80% were Black women).

Enzyme-linked immunosorbent assay

ELISAs were completed using plates coated with the RBD of the SARS-CoV-2 spike protein using a previously described protocol with slight modifications (6, 25). Plasmids for expressing this protein were provided by F. Krammer (Mt. Sinai). SARS-CoV-2 RBD proteins were produced in 293F cells and purified using nickel–nitrilotriacetic acid (Ni-NTA) resin (Qiagen). The supernatant was incubated for 2 hours with Ni-NTA resin at room temperature before the Ni-NTA resin was collected using gravity flow columns, and the protein was eluted. After buffer exchange into phosphate-buffered saline (PBS), the purified protein was stored in aliquots at -80°C . ELISA plates (Immulon 4 HBX, Thermo Fisher Scientific) were coated overnight at 4°C with PBS (50 μl per well) or a recombinant protein (2 $\mu\text{g}/\text{ml}$) diluted in PBS. The next day, ELISA plates were washed three times with PBS containing 0.1% Tween 20 (PBS-T) and blocked for 1 hour with PBS-T supplemented with 3% nonfat milk powder. Before testing in ELISA, serum samples were heat-inactivated at 56°C for 1 hour. Serum samples were serially diluted in twofold in 96-well round-bottom plates in PBS-T supplemented with 1% nonfat milk powder (dilution buffer), starting at a 1:50 dilution. Next, ELISA plates were washed three times with PBS-T, and 50 μl of serum dilution was added to each well. Plates were incubated for 2 hours at room temperature using a plate mixer. Plates were washed again three times with PBS-T before 50 μl of horseradish peroxidase (HRP)–labeled goat anti-human IgG (1:5000; Jackson ImmunoResearch Laboratories) or goat anti-human IgM-HRP (1:1000; SouthernBiotech) secondary antibodies were added. After 1-hour incubation at room temperature using a plate mixer, plates were washed three times with PBS-T, and 50 μl of SureBlue 3,3',5,5'-tetramethylbenzidine substrate (KPL) was added to each well. Five minutes later, 25 μl of 250 mM hydrochloric acid was added to each well to stop the reaction. Plates were read at an optical density (OD) of 450 nm using the SpectraMax 190 microplate reader (Molecular Devices). Background OD values from the plates coated with PBS were subtracted from the OD values from plates coated with recombinant protein. A dilution series of the IgG monoclonal antibody CR3022, which is reactive to the SARS-CoV-2 spike protein, was included on each plate as a control to adjust for interassay variability. The IgG CR3022 monoclonal antibody was included on both IgG and IgM plates, and an anti-human

IgG-HRP secondary antibody was added to these standardization wells on both IgG and IgM plates. In essence, the CR3022 monoclonal antibody was used to set the OD threshold on each plate and to ensure that the same OD threshold was used on all plates, including both IgG and IgM assays. Serum antibody concentrations were reported as arbitrary units relative to the CR3022 monoclonal antibody. Plasmids to express the CR3022 monoclonal antibody were provided by I. Wilson (Scripps). All samples were first tested in duplicate at a 1:50 serum dilution. Samples with an IgG and/or IgM concentration above the lower limit of detection (0.20 arbitrary units) were repeated in at least a seven-point dilution series to obtain quantitative results.

Establishment of an ELISA cutoff to distinguish seropositive versus seronegative

We used results from the 2019 cohort (Fig. 2A) to set ELISA cutoffs for seropositivity and seronegativity. Over the course of establishing our serological assay, we identified rare individuals who had pre-pandemic SARS-CoV-2 cross-reactive serum antibodies. Most of these individuals had very low levels of cross-reactive SARS-CoV-2 antibodies. We found that ~1% of samples from the pre-pandemic 2019 cohort had IgG and/or IgM levels of >0.48 arbitrary units, which was subsequently used as the cutoff for defining seropositivity in the 2020 cohort.

Statistical methods

Standard descriptive analyses using χ^2 test, Fisher's exact test, and Mann-Whitney *U* test as appropriate, compared the demographic and clinical characteristics between the seropositive and seronegative women. CIs for proportions were computed using the Clopper-Pearson (exact) method. Statistical significance was set at $P < 0.05$. Statistical analyses were performed using Stata version 16 (StataCorp, College Station, TX) and Prism version 8 (GraphPad Software). Figure 1 was created using QGIS version 3.12.3.

SUPPLEMENTARY MATERIALS

immunology.sciencemag.org/cgi/content/full/5/49/eabd5709/DC1

Table S1. Relative levels of SARS-CoV-2 IgG and IgM in serum collected from seropositive pregnant women ($n = 80$).

[View/request a protocol for this paper from Bio-protocol.](#)

REFERENCES AND NOTES

- D. Wang, B. Hu, C. Hu, F. Zhu, X. Liu, J. Zhang, B. Wang, H. Xiang, Z. Cheng, Y. Xiong, Y. Zhao, Y. Li, X. Wang, Z. Peng, Clinical Characteristics of 138 Hospitalized Patients with 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA* **323**, 1061–1069 (2020).
- J. Abbasi, The promise and peril of antibody testing for COVID-19. *JAMA* 10.1001/jama.2020.6170, (2020).
- D. Crook, Evaluation of antibody testing for SARS-CoV-2 using ELISA and lateral flow immunoassays. *medRxiv* 10.1101/2020.04.15.20066407, (2020).
- J. D. Whitman, J. Hiatt, C. T. Mowery, B. R. Shy, R. Yu, T. N. Yamamoto, U. Rathore, G. M. Goldgof, C. Whitty, J. M. Woo, A. E. Gallman, T. E. Miller, A. G. Levine, D. N. Nguyen, S. P. Bapat, J. Balcerak, S. A. Bylisma, A. M. Lyons, S. Li, A. W.-Y. Wong, E. M. Gillis-Buck, Z. B. Steinhart, Y. Lee, R. Apathy, M. J. Lipke, J. A. Smith, T. Zheng, I. C. Boothby, E. Isaza, J. Chan, D. D. Acenas II, J. Lee, T. A. Macrae, T. S. Kyaw, D. Wu, D. L. Ng, W. G. V. A. York, H. A. Eskandarian, P. C. Callaway, L. Warriar, M. E. Moreno, J. Levan, L. Torres, L. A. Farrington, R. Loudermilk, K. Koshal, K. C. Zorn, W. F. Garcia-Beltran, D. Yang, M. G. Astudillo, B. E. Bernstein, J. A. Gelfand, E. T. Ryan, R. C. Charles, A. J. Iafraite, J. G. Lennnerz, S. Miller, C. Y. Chiu, S. L. Stramer, M. R. Wilson, A. Manglik, C. J. Ye, N. J. Krogan, M. S. Anderson, J. G. Cyster, J. D. Ernst, A. H. B. Wu, K. L. Lynch, C. Bern, P. D. Hsu, A. Marson, Test performance evaluation of SARS-CoV-2 serological assays. *medRxiv* 2020.04.25.20074856, (2020).
- Vital Statistics Annual Report; www.health.pa.gov/topics/HealthStatistics/VitalStatistics/PAVitalStatistics/Pages/vital-statistics-report.aspx.
- F. Amanat, D. Stadlbauer, S. Strohmaier, T. H. O. Nguyen, V. Chromikova, M. McMahon, K. Jiang, G. A. Arunkumar, D. Jurczyszczak, J. Polanco, M. Bernudez-Gonzalez, G. Kleiner, T. Aydllo, L. Miorin, D. Fierer, L. A. Lugo, E. M. Kojic, J. Stoeber, S. T. H. Liu, C. Cunningham-Rundles, P. L. Felgner, T. Moran, A. Garcia-Sastre, D. Caplivski, A. Cheng, K. Kedzierska, O. Vapalahti, J. M. Hepojoki, V. Simon, F. Krammer, A serological assay to detect SARS-CoV-2 seroconversion in humans. *Nat. Med.* **26**, 1033–1036 (2020).
- L. Kuri-Cervantes, M. B. Pampena, W. Meng, A. M. Rosenfeld, C. A. G. Ittner, A. R. Weisman, R. S. Agyekum, D. Mathew, A. E. Baxter, L. A. Vella, O. Kuthuru, S. A. Apostolidis, L. Bershaw, J. Dougherty, A. R. Greenplate, A. Pattekar, J. Kim, N. Han, S. Gouma, M. E. Weirick, C. P. Arevalo, M. J. Bolton, E. C. Goodwin, E. M. Anderson, S. A. Apostolidis, T. K. Jones, N. S. Mangalmurti, E. T. L. Prak, E. J. Wherry, N. J. Meyer, M. R. Betts, Comprehensive mapping of immune perturbations associated with severe COVID-19. *Sci. Immunol.* **5**, eabd7114 (2020).
- D. Mathew, J. R. Giles, A. E. Baxter, D. A. Oldridge, A. R. Greenplate, J. E. Wu, C. Alanio, L. Kuri-Cervantes, M. B. Pampena, K. D'Andrea, S. Manne, Z. Chen, Y. J. Huang, J. P. Reilly, A. R. Weisman, C. A. G. Ittner, O. Kuthuru, J. Dougherty, K. Nzingha, N. Han, J. Kim, A. Pattekar, E. C. Goodwin, E. M. Anderson, M. E. Weirick, S. Gouma, C. P. Arevalo, M. J. Bolton, F. Chen, S. F. Lacey, H. Ramage, S. Cherry, S. E. Hensley, S. A. Apostolidis, A. C. Huang, L. A. Vella; The UPenn COVID Processing Unit, M. R. Betts, N. J. Meyer, E. J. Wherry, Deep immune profiling of COVID-19 patients reveals distinct immunotypes with therapeutic implications. *Science*, eabc8511 (2020).
- City Provides Update on COVID-19 for Wednesday, June 3, 2020 (Department of Public Health, City of Philadelphia, 2020); www.phila.gov/2020-06-03-city-provides-update-on-covid-19-for-wednesday-june-3-2020/.
- F. Zeng, C. Dai, P. Cai, J. Wang, L. Xu, J. Li, G. Hu, L. Wang, A comparison study of SARS-CoV-2 IgG antibody between male and female COVID-19 patients: A possible reason underlying different outcome between gender. *medRxiv* 2020.03.26.20040709, (2020).
- D. Devakumar, S. Selvarajah, G. Shannon, K. Muraya, S. Lasoye, S. Corona, Y. Paradies, I. Abubakar, E. T. Achiume, Racism, the public health crisis we can no longer ignore. *Lancet* **395**, e112–e113 (2020).
- D. Devakumar, S. S. Bhopal, G. Shannon, COVID-19: The great unequaliser. *J. R. Soc. Med.* **113**, 234–235 (2020).
- E. Abuelgasim, L. J. Saw, M. Shirke, M. Zeinah, A. Harky, COVID-19: Unique public health issues facing Black, Asian and minority ethnic communities. *Curr. Probl. Cardiol.* **45**, 100621 (2020).
- H. Chen, J. Guo, C. Wang, F. Luo, X. Yu, W. Zhang, J. Li, D. Zhao, D. Xu, Q. Gong, J. Liao, H. Yang, W. Hou, Y. Zhang, Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: A retrospective review of medical records. *Lancet* **395**, 809–815 (2020).
- R. Chen, Y. Zhang, L. Huang, B.-H. Cheng, Z.-Y. Xia, Q.-T. Meng, Safety and efficacy of different anesthetic regimens for parturients with COVID-19 undergoing Cesarean delivery: A case series of 17 patients. *Can. J. Anaesth.* **67**, 655–663 (2020).
- N. Li, L. Han, M. Peng, Y. Lv, Y. Ouyang, K. Liu, L. Yue, Q. Li, G. Sun, L. Chen, L. Yang, Maternal and neonatal outcomes of pregnant women with COVID-19 pneumonia: A case-control study. *medRxiv* 2020.03.10.20033605, (2020).
- D. Liu, L. Li, X. Wu, D. Zheng, J. Wang, L. Yang, C. Zheng, Pregnancy and perinatal outcomes of women with coronavirus disease (COVID-19) Pneumonia: A preliminary analysis. *AJR Am. J. Roentgenol.* **215**, 127–132 (2020).
- W. Liu, Q. Wang, Q. Zhang, L. Chen, J. Chen, B. M. Zhang, S. Wang, L. Xia, L. Huang, K. B. Wang, L. B. Liang, Y. M. Zhang, L. Turtle, D. Lissauer, K. Lan, L. Feng, Y. Liu, Z. Sun, *Coronavirus Disease 2019 (COVID-19) During Pregnancy: A Case Series* (Preprints, 2020); www.preprints.org.
- X. Wang, Z. Zhou, J. Zhang, F. Zhu, Y. Tang, X. Shen, A case of 2019 Novel Coronavirus in a pregnant woman with preterm delivery. *Clin. Infect. Dis.* **71**, 844–846 (2020).
- L. Zhang, Y. Jiang, M. Wei, B. H. Cheng, X. C. Zhou, J. Li, J. H. Tian, L. Dong, R. H. Hu, Analysis of the pregnancy outcomes in pregnant women with COVID-19 in Hubei Province. *Zhonghua Fu Chan Ke Za Zhi* **55**, 166–171 (2020).
- H. Zhu, L. Wang, C. Fang, S. Peng, L. Zhang, G. Chang, S. Xia, W. Zhou, Clinical analysis of 10 neonates born to mothers with 2019-nCoV pneumonia. *Transl. Pediatr.* **9**, 51–60 (2020).
- M. Knight, K. Bunch, N. Vousden, E. Morris, N. Simpson, C. Gale, P. O'Brien, M. Quigley, P. Brocklehurst, J. J. Kurinczuk; UK Obstetric Surveillance System SARS-CoV-2 Infection in Pregnancy Collaborative Group, Characteristics and outcomes of pregnant women admitted to hospital with confirmed SARS-CoV-2 infection in UK: National population based cohort study. *BMJ* **369**, m2107 (2020).
- H. Zeng, C. Xu, J. Fan, Y. Tang, Q. Deng, W. Zhang, X. Long, Antibodies in Infants Born to Mothers with COVID-19 Pneumonia. *JAMA* **323**, 1848–1849 (2020).
- M. C. Alzamora, T. Paredes, D. Caceres, C. M. Webb, L. M. Valdez, M. La Rosa, Severe COVID-19 during pregnancy and possible vertical transmission. *Am. J. Perinatol.* **37**, 861–865 (2020).

25. D. Stadlbauer, F. Amanat, V. Chromikova, K. Jiang, S. Strohmeier, G. A. Arunkumar, J. Tan, D. Bhavsar, C. Capuano, E. Kirkpatrick, P. Meade, R. N. Brito, C. Teo, M. McMahon, V. Simon, F. Krammer, SARS-CoV-2 Seroconversion in Humans: A detailed protocol for a serological assay, antigen production, and test setup. *Curr. Protoc. Microbiol.* **57**, e100 (2020).

Acknowledgments: We thank all members of the Wherry Lab and the Penn COVID-19 Sample Processing Unit for sample procurement, processing, and logistics. We thank the staff of the PMBB. We thank F. Krammer (Mt. Sinai) for sending us the SARS-CoV-2 spike RBD expression plasmids. We thank S. Melly (Drexel University) for the assistance in geographic analyses.

Funding: This work was supported by institutional funds from the University of Pennsylvania and NIH grants AI082630 (to E.J.W.) and UL1TR001878 (to D.J.R.). We thank J. Lurie, J. Embiid, J. Harris, and D. Blitzer for philanthropic support. E.J.W. is supported by the Parker Institute for Cancer Immunotherapy, which supports the cancer immunology program at University of Pennsylvania. **Author contributions:** D.D.F. conceptualized and designed the study, collected the data, drafted the initial manuscript, and revised the manuscript. S.G. led the serological experiments, collected the data, and revised the manuscript. M.B.D. designed the data collection instruments, collected the data, carried out the analyses, and revised the manuscript. S.M. conceptualized and designed the study, designed the data collection instruments, carried out the analyses, and revised the manuscript. M.R.P. and E.C.W. collected data and revised the manuscript. J.S.G. conceptualized and designed the study and revised the manuscript. C.P.A., M.J.B., M.E.W., E.C.G., and E.M.A. completed serological assays, analyzed the data, and revised the manuscript. A.R.G., J.K., N.H., A.P., and J.D. obtained and proceeded samples from recovered donors. O.K. and L.A.V. designed and established recovered donor cohort. D.M. processed and characterized samples from recovered donors. A.E.B. oversaw acquisition, processing, and characterization of samples from recovered donors. J.W. supervised the recruitment of participants in PMBB and identification of samples for serology

testing. A.V. analyzed demographic data of PMBB participants. R.L. provided samples for the pre-pandemic pregnant controls. J.S.M. provided statistical advice, performed statistical analyses, and revised the paper. D.J.R. provided input on the use of PMBB controls and revised the manuscript. M.A.E. provided input and samples for the pre-pandemic pregnant controls and revised the manuscript. E.J.W. designed, established, and oversaw healthy donor cohort studies and made revisions to the manuscript. K.M.P. conceptualized and designed the study, coordinated and supervised data collection, and revised the manuscript. S.E.H. conceptualized and designed the study, coordinated and supervised serological studies, and revised the manuscript. **Competing interests:** S.E.H. has received consultancy fee from Sanofi Pasteur, Lumen, Novavax, and Merck for work unrelated to this report. E.J.W. is a member of the Parker Institute for Cancer Immunotherapy. E.J.W. has consulting agreements with and/or is on the scientific advisory board for Merck, Roche, Pieris, Elstar, and Surface Oncology. E.J.W. is a founder of Surface Oncology and Arsenal Biosciences. E.J.W. has a patent licensing agreement on the PD-1 pathway with Roche/Genentech. All other authors declare that they have no competing interests. **Data and materials availability:** All data needed to evaluate the conclusions in the paper are present in the paper or the Supplementary Materials.

Submitted 29 June 2020

Accepted 24 July 2020

Published 29 July 2020

Final published 24 August 2020

10.1126/sciimmunol.abd5709

Citation: D. D. Flannery, S. Gouma, M. B. Dhudasia, S. Mukhopadhyay, M. R. Pfeifer, E. C. Woodford, J. S. Gerber, C. P. Arevalo, M. J. Bolton, M. E. Weirick, E. C. Goodwin, E. M. Anderson, A. R. Greenplate, J. Kim, N. Han, A. Pattekar, J. Dougherty, O. Kuthuru, D. Mathew, A. E. Baxter, L. A. Vella, J. Weaver, A. Verma, R. Leite, J. S. Morris, D. J. Rader, M. A. Elovitz, E. J. Wherry, K. M. Puopolo, S. E. Hensley, SARS-CoV-2 seroprevalence among parturient women in Philadelphia. *Sci. Immunol.* **5**, eabd5709 (2020).

SARS-CoV-2 seroprevalence among parturient women in Philadelphia

Dustin D. Flannery, Sigrid Gouma, Miren B. Dhudasia, Sagori Mukhopadhyay, Madeline R. Pfeifer, Emily C. Woodford, Jeffrey S. Gerber, Claudia P. Arevalo, Marcus J. Bolton, Madison E. Weirick, Eileen C. Goodwin, Elizabeth M. Anderson, Allison R. Greenplate, Justin Kim, Nicholas Han, Ajinkya Pattekar, Jeanette Dougherty, Oliva Kuthuru, Divij Mathew, Amy E. Baxter, Laura A. Vella, JoEllen Weaver, Anurag Verma, Rita Leite, Jeffrey S. Morris, Daniel J. Rader, Michal A. Elovitz, E. John Wherry, Karen M. Puopolo and Scott E. Hensley

Sci. Immunol. **5**, eabd5709.
First published 29 July 2020
DOI: 10.1126/sciimmunol.abd5709

Monitoring SARS-CoV-2 exposure

When the SARS-CoV-2 pandemic started, stay-at-home orders led to postponement of routine medical appointments, making it difficult to perform community-level seroprevalence studies. Pregnant women, particularly those near their delivery dates, continued to have regular interactions with their medical care providers. Flannery *et al.* measured the exposure rate of SARS-CoV-2 in parturient women in Philadelphia who came to the hospital to deliver between 4 April and 3 June 2020. Using IgG and IgM antibody tests, they evaluated exposure of 1293 parturient women to SARS-CoV-2 and found that 6.2% of these women had evidence of exposure to the virus. They found considerable racial differences in exposure rates, with higher rates of seropositivity in Black/non-Hispanic and Hispanic/Latino women as compared with Asian and White/non-Hispanic women.

ARTICLE TOOLS

<http://immunology.sciencemag.org/content/5/49/eabd5709>

SUPPLEMENTARY MATERIALS

<http://immunology.sciencemag.org/content/suppl/2020/07/27/5.49.eabd5709.DC1>

REFERENCES

This article cites 19 articles, 4 of which you can access for free
<http://immunology.sciencemag.org/content/5/49/eabd5709#BIBL>

Use of this article is subject to the [Terms of Service](#)

Science Immunology (ISSN 2470-9468) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. The title *Science Immunology* is a registered trademark of AAAS.

Copyright © 2020 The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. Distributed under a Creative Commons Attribution License 4.0 (CC BY).